



DEPARTMENT: AGRICULTURE

Lungsickness

**Lungsickness or contagious bovine pleuropneumonia (CBPP)
is a disease of cattle in Africa**



- △ The disease is present in cattle of our neighbouring countries, Namibia and Botswana.
- △ The symptoms of lungsickness should be recognised by every farmer to prevent the spreading of the disease.

How the disease is transmitted

- △ The bacteria causing lungsickness is transmitted when infected cattle come into close contact with healthy cattle.
- △ This often happens at watering places or when cattle are kraaled at night.

Symptoms of lungsickness

- △ This disease affects the lungs and chest of cattle and causes:
 - fever
 - coughing
 - listlessness
 - difficulty in breathing
 - discharges from the nose and
 - loss of condition.

**Contact your nearest
state veterinarian if
animals in your herd
have these symptoms**

How to prevent lungsickness in your herd

- △ Inspect cattle regularly for symptoms of the disease.
- △ Buy only healthy cattle.
- △ If possible do not allow your cattle to come into contact with strange cattle.

Controlling lungsickness

- △ Herds with infected cattle may have to be slaughtered to control the disease.
- △ Cattle belonging to neighbours must be vaccinated.
- △ Movement of cattle will not be allowed.
- △ Cattle farmers must be able to recognise the symptoms of lungsickness.

For further information, contact the Directorate Animal Health
Private Bag X138, Pretoria 0001
Tel: (012) 319 7679

or

Resource Centre, Department of Agriculture
Tel: (012) 319 7141/7085

This publication is available on the web: www.nda.agric.za/publications

2002 Revised edition

Compiled by the Department of Agriculture, Directorate Agricultural Information Services

Printed and published in the Republic of South Africa by the Department of Agriculture and obtainable from the Resource Centre, Directorate Agricultural Information Services
Private Bag X144, Pretoria, 0001 South Africa