




Sexually transmitted infection

From Wikipedia, the free encyclopedia

Sexually transmitted infections (STI), also referred to as **sexually transmitted diseases (STD)** and **venereal diseases (VD)**, are infections that are commonly spread by sex, especially vaginal intercourse, anal sex and oral sex. Most STIs initially do not cause symptoms.^[1] This results in a greater risk of passing the disease on to others.^{[2][3]} Symptoms and signs of disease may include vaginal discharge, penile discharge, ulcers on or around the genitals, and pelvic pain. STIs acquired before or during birth may result in poor outcomes for the baby. Some STIs may cause problems with the ability to get pregnant.^[1]

More than 30 different bacteria, viruses, and parasites can cause STIs.^[1] Bacterial STIs include chlamydia, gonorrhea, and syphilis among others. Viral STIs include genital herpes, HIV/AIDS, and genital warts among others. Parasitic STIs include trichomoniasis among others. While usually spread by sex, some STIs can also be spread by non-sexual contact with contaminated blood and tissues, breastfeeding, or during childbirth.^[1] STI diagnostic tests are easily available in the

Sexually transmitted infection



"Syphilis is a dangerous disease, but it can be cured." Poster encouraging treatment showing text and design of an anchor and a cross. Published between 1936 and 1938.

Classification and external resources

Specialty	Infectious disease
ICD-10	A64 (http://apps.who.int/classifications/icd10/browse/2016/en#/A64)
ICD-9-CM	099.9 (http://www.icd9data.com/getICD9Code.aspx?icd9=099.9)
DiseasesDB	27130 (http://www.diseasesdatabase.com/ddb27130.htm)
Patient UK	Sexually transmitted infection (http://patient.info/doctor/sexually-transmitted-infections-pro)

developed world, but this is often not the case in the developing world.^[1]

MeSH D012749 (https://www.nlm.nih.gov/cgi/mesh/2016/MB_cgi?field=uid&term=D012749)

The most effective way of preventing STIs is by not having sex.^[4] Some vaccinations may also decrease the risk of certain infections including hepatitis B and some types of HPV.^[4] Safer sex practices such as use of condoms, having a smaller number of sexual partners, and being in a relationship where each person only has sex with the other also decreases the risk.^{[1][4]} Circumcision in males may be effective to prevent some infections.^[1] Most STIs are treatable or curable.^[1] Of the most common infections, syphilis, gonorrhea, chlamydia, trichomoniasis are curable, while herpes, hepatitis B, HIV/AIDS, and HPV are treatable but not curable.^[1] Resistance to certain antibiotics is developing among some organisms such as gonorrhea.^[5]

In 2008, it was estimated that 500 million people were infected with either syphilis, gonorrhea, chlamydia or trichomoniasis.^[1] At least an additional 530 million people have genital herpes and 290 million women have human papillomavirus.^[1] STIs other than HIV resulted in 142,000 deaths in 2013.^[6] In the United States there were 19 million new cases of sexually transmitted infections in 2010.^[7] Historical documentation of STIs date back to at least the Ebers papyrus around 1550 BC and the Old Testament.^[8] There is often shame and stigma associated with these infections.^[1] The term *sexually transmitted infection* is generally preferred over the terms *sexually transmitted disease* and *venereal disease*, as it includes those who do not have symptomatic disease.^[9]

Contents

- 1 Classification
- 2 Signs and symptoms
- 3 Cause
 - 3.1 Transmission
 - 3.2 Bacterial
 - 3.3 Fungal
 - 3.4 Viral
 - 3.5 Parasites
 - 3.6 Protozoal
 - 3.7 Main types
 - 3.8 Unscreened
- 4 Pathophysiology
- 5 Prevention
 - 5.1 Vaccines
 - 5.2 Condoms
 - 5.3 Nonoxynol-9
- 6 Screening
- 7 Diagnosis

- 8 Management
- 9 Epidemiology
- 10 History
- 11 Other names or slang terms
- 12 Plants
- 13 References
- 14 Further reading
- 15 External links

Classification

Until the 1990s, STIs were commonly known as *venereal diseases*, the word *venereal* being derived from the Latin word *venereus*, and meaning relating to sexual intercourse or desire, ultimately derived from Venus, the Roman goddess of love.^[10] "Social disease" was a phrase used as a euphemism.

The World Health Organization (WHO) has recommended *sexually transmitted infection* as the preferred term since 1999.^[9] *Sexually transmitted infection* is a broader term than *sexually transmitted disease*.^[11] An infection is a colonization by a parasitic species, which may not cause any adverse effects. In a disease, the infection leads to impaired or abnormal function. In either case, the condition may not exhibit signs or symptoms. Increased understanding of infections like HPV, which infects a significant portion of sexually active individuals but cause disease in only a few, has led to increased use of the term *STI*. Public health officials originally introduced the term *sexually transmitted infection*, which clinicians are increasingly using alongside the term *sexually transmitted disease* in order to distinguish it from the former.^[12]

STD may refer only to infections that are causing diseases, or it may be used more loosely as a synonym for *STI*. Most of the time, people do not know that they are infected with an STI until they are tested or start showing symptoms of disease. Moreover, the term *sexually transmissible disease* is sometimes used since it is less restrictive in consideration of other factors or means of transmission. For instance, meningitis is transmissible by means of sexual contact but is not labeled an STI because sexual contact is not the primary vector for the pathogens that cause meningitis. This discrepancy is addressed by the probability of infection by means *other than sexual contact*. In general, an STI is an infection that has a negligible probability of transmission by means other than sexual contact, but has a realistic means of transmission by sexual contact (more sophisticated means—blood transfusion, sharing of hypodermic needles—are not taken into account). Thus, one may presume that, if a person is infected with an STI, e.g., chlamydia, gonorrhea, genital herpes, HPV it was transmitted to him/her by means of sexual contact.



A poster from the Office for Emergency Management. Office of War Information, 1941-1945

Signs and symptoms

Not all STIs are symptomatic, and symptoms may not appear immediately after infection. In some instances a disease can be carried with no symptoms, which leaves a greater risk of passing the disease on to others. Depending on the disease, some untreated STIs can lead to infertility, chronic pain or even death.^[13]

The presence of an STI in prepubescent children may indicate sexual abuse.^[14]

Cause

Transmission

The risks and transmission probabilities of sexually transmitted diseases are summarized by act in the table:^{[15][16][17][18][19][20][21][22][23][24][25][26][27]}

Risk per unprotected sexual act with an infected person		
	Known risks	Possible
Performing oral sex on a man	<ul style="list-style-type: none"> ▪ Throat chlamydia^[15] ▪ Throat gonorrhea^[15] (25–30%) ▪ Herpes (rare) ▪ HPV^[16] ▪ Syphilis^[15] (1%)^[17] 	<ul style="list-style-type: none"> ▪ Hepatitis B (low risk)^[18] ▪ HIV (0.01%)^[19] ▪ Hepatitis C (unknown)
Performing oral sex on a woman	<ul style="list-style-type: none"> ▪ Herpes ▪ HPV^[16] 	<ul style="list-style-type: none"> ▪ Throat gonorrhea^[15] ▪ Throat chlamydia^[15]
Receiving oral sex—man	<ul style="list-style-type: none"> ▪ Chlamydia ▪ Gonorrhea^[15] ▪ Herpes ▪ Syphilis^[15] (1%)^[17] 	<ul style="list-style-type: none"> ▪ HPV
Receiving oral sex—woman	<ul style="list-style-type: none"> ▪ Herpes 	<ul style="list-style-type: none"> ▪ HPV ▪ Bacterial Vaginosis^[15] ▪ Gonorrhea^[15]
Vaginal sex—man	<ul style="list-style-type: none"> ▪ Chlamydia (30–50%)^[18] ▪ Crabs ▪ Scabies ▪ Gonorrhea (22%)^[20] ▪ Hepatitis B ▪ Herpes (0.07% for HSV-2)^[21] ▪ HIV (0.05%)^{[19][21]} ▪ HPV (high: around 40-50%)^[22] ▪ Mycoplasma hominis infection^{[28][29][30][31]} ▪ Syphilis ▪ Trichomoniasis ▪ Ureaplasma infection^{[32][33]} 	<ul style="list-style-type: none"> ▪ Hepatitis C
Vaginal sex—woman	<ul style="list-style-type: none"> ▪ Chlamydia (30–50%)^[18] ▪ Crabs ▪ Scabies 	<ul style="list-style-type: none"> ▪ Hepatitis C

	<ul style="list-style-type: none"> ▪ Gonorrhea (47%)^[23] ▪ Hepatitis B (50–70%) ▪ Herpes ▪ HIV (0.1%)^[19] ▪ HPV (high;^[18] around 40-50%)^[22] ▪ Mycoplasma hominis infection^{[28][29][31]} ▪ Syphilis ▪ Trichomoniasis ▪ Ureaplasma infection^{[32][33]} 	
Anal sex—insertive	<ul style="list-style-type: none"> ▪ Chlamydia ▪ Crabs ▪ Scabies (40%) ▪ Gonorrhea ▪ Hepatitis B ▪ Herpes ▪ HIV (0.62%)^[25] ▪ HPV ▪ Syphilis (14%)^[17] 	<ul style="list-style-type: none"> ▪ Hepatitis C
Anal sex—receptive	<ul style="list-style-type: none"> ▪ Chlamydia ▪ Crabs ▪ Scabies ▪ Gonorrhea ▪ Hepatitis B ▪ Herpes ▪ HIV (1.7%)^[25] ▪ HPV ▪ Syphilis (1.4%)^[17] 	<ul style="list-style-type: none"> ▪ Hepatitis C
Anilingus	<ul style="list-style-type: none"> ▪ Amebiasis ▪ Cryptosporidiosis (1%) ▪ Giardiasis^[34] ▪ Hepatitis A^[35] (1%) ▪ Shigellosis^[36] (1%) 	<ul style="list-style-type: none"> ▪ HPV (1%)

Bacterial

- Chancroid (*Haemophilus ducreyi*)
- Chlamydia (*Chlamydia trachomatis*)


- Gonorrhea (*Neisseria gonorrhoeae*), colloquially known as "the clap"
- Granuloma inguinale or (*Klebsiella granulomatis*)
- Mycoplasma genitalium^{[37][38][39]}
- Mycoplasma hominis^{[28][29][30][31][40]}
- Syphilis (*Treponema pallidum*)
- Ureaplasma infection^{[32][33]}

Fungal

- Candidiasis (yeast infection)

Viral

- Viral hepatitis (Hepatitis B virus)—saliva, venereal fluids. (Note: Hepatitis A and Hepatitis E are transmitted via the fecal-oral route; Hepatitis C is rarely sexually transmittable,^[41] and the route of transmission of Hepatitis D (only if infected with B) is uncertain, but may include sexual transmission.^{[42][43][44]})
- Herpes simplex (Herpes simplex virus 1, 2) skin and mucosal, transmissible with or without visible blisters
- HIV (*Human Immunodeficiency Virus*)—venereal fluids, semen, breast milk, blood
- HPV (*Human Papillomavirus*)—skin and mucosal contact. 'High risk' types of HPV cause almost all cervical cancers, as well as some anal, penile, and vulvar cancer. Some other types of HPV cause genital warts.
- Molluscum contagiosum (molluscum contagiosum virus MCV)—close contact



Micrograph showing the viral cytopathic effect of herpes (ground glass nuclear inclusions, multi-nucleation). Pap test. Pap stain.

Parasites

- Crab louse, colloquially known as "crabs" or "pubic lice" (*Pthirus pubis*)
- Scabies (*Sarcoptes scabiei*)

Protozoal

- Trichomoniasis (*Trichomonas vaginalis*), colloquially known as "trich"

Main types

Sexually transmitted infections include:


- Chlamydia is a sexually transmitted infection caused by the bacterium *Chlamydia trachomatis*. In women, symptoms may include abnormal vaginal discharge, burning during urination, and bleeding in between periods, although most women do not experience any symptoms.^[45]

Symptoms in men include pain when urinating, and abnormal discharge from their penis.^[46] If left untreated in both men and women, Chlamydia can infect the urinary tract and potentially lead to pelvic inflammatory disease (PID). PID can cause serious problems during pregnancy and even has the potential to cause infertility. It can cause a woman to have a potentially deadly ectopic pregnancy, in which the child is born outside of the uterus. However, Chlamydia can be cured with antibiotics.

- The two most common forms of herpes are caused by infection with herpes simplex virus (HSV). HSV-1 is typically acquired orally and causes cold sores, HSV-2 is usually acquired during sexual contact and affects the genitals, however either strain may affect either site.^[47] Some people are asymptomatic or have very mild symptoms. Those that do experience symptoms usually notice them 2 to 20 days after exposure which last 2 to 4 weeks. Symptoms can include small fluid-filled blisters, headaches, backaches, itching or tingling sensations in the genital or anal area, pain during urination, Flu like symptoms, swollen glands, or fever. Herpes is spread through skin contact with a person infected with the virus. The virus affects the areas where it entered the body. This can occur through kissing, vaginal intercourse, oral sex or anal sex. The virus is most infectious during times when there are visible symptoms, however those who are asymptomatic can still spread the virus through skin contact.^[48] The primary attack is the most severe because the body does not have any antibodies built up. After the primary attack, one might have recurring attacks that are milder or might not even have future attacks. There is no cure for the disease but there are antiviral medications that treat its symptoms and lower the risk of transmission (Valtrex). Although HSV-1 is typically the "oral" version of the virus, and HSV-2 is typically the "genital" version of the virus, a person with HSV-1 orally CAN transmit that virus to their partner genitally. The virus, either type, will settle into a nerve bundle either at the top of the spine, producing the "oral" outbreak, or a second nerve bundle at the base of the spine, producing the genital outbreak.
- The human papillomavirus (HPV) is the most common STI in the United States.^[49] There are more than 40 different strands of HPV and many do not cause any health problems. In 90% of cases the body's immune system clears the infection naturally within 2 years.^[50] Some cases may not be cleared and can lead to genital warts (bumps around the genitals that can be small or large, raised or flat, or shaped like cauliflower) or cervical cancer and other HPV related cancers. Symptoms might not show up until advanced stages. It is important for women to get pap smears in order to check for and treat cancers. There are also two vaccines available for women (Cervarix and Gardasil) that protect against the types of HPV that cause cervical cancer. HPV can be passed through genital-to-genital contact as well as during oral sex. It is important to remember that the infected partner might not have any symptoms.
- Gonorrhea is caused by bacterium that lives on moist mucous membranes in the urethra, vagina, rectum, mouth, throat, and eyes. The infection can spread through contact with the penis, vagina, mouth or anus. Symptoms of Gonorrhea usually appear 2 to 5 days after contact with an infected partner however, some men might not notice symptoms for up to a month. Symptoms in men include burning and pain while urinating, increased urinary frequency, discharge from the penis (white, green, or yellow in color), red or swollen urethra, swollen or tender testicles, or sore throat. Symptoms in women may include vaginal discharge, burning or itching while urinating, painful sexual intercourse, severe pain in lower abdomen (if infection spreads to fallopian tubes), or fever (if infection spreads to fallopian tubes), however many women do not show any symptoms.^[51] There are some antibiotic resistant strains for Gonorrhea but most cases can be cured with antibiotics.
- Syphilis is an STI caused by a bacterium. Untreated, it can lead to complications and death.^[52] Clinical manifestations of syphilis include the ulceration of the uro-genital tract, mouth or rectum;

if left untreated the symptoms worsen. In recent years, the prevalence of syphilis has declined in Western Europe, but it has increased in Eastern Europe (former Soviet states). A high incidence of syphilis can be found in places such as Cameroon, Cambodia, Papua New Guinea.^[53] Syphilis infections are increasing in the United States.^[54]

- Trichomoniasis is a common STI that is caused by infection with a protozoan parasite called *Trichomonas vaginalis*.^[55] Trichomoniasis affects both women and men, but symptoms are more common in women.^[56] Most patients are treated with an antibiotic called metronidazole, which is very effective.^[57]
- HIV (human immunodeficiency virus) damages the body's immune system, which interferes with its ability to fight off disease-causing agents. The virus kills CD4 cells, which are white blood cells that help fight off various infections. HIV is carried in body fluids, and is spread by sexual activity. It can also be spread by contact with infected blood, breast feeding, childbirth, and from mother to child during pregnancy.^[58] When HIV is at its most advanced stage, an individual is said to have AIDS (acquired immunodeficiency syndrome).^[59] There are different stages of the progression of and HIV infection. The stages include primary infection, asymptomatic infection, symptomatic infection, and AIDS. In the primary infection stage, an individual will have flu like symptoms (headache, fatigue, fever, muscle aches) for about 2 weeks. In the asymptomatic stage, symptoms usually disappear, and the patient can remain asymptomatic for years. When HIV progresses to the symptomatic stage, the immune system is weakened, and has a low cell count of CD4+ T Cells. When the HIV infection becomes life-threatening, it is called AIDS. People with AIDS fall prey to opportunistic infections and die as a result.^[45] When the disease was first discovered in the 1980s, those who had AIDS were not likely to live longer than a few years. There are now antiretroviral drugs (ARVs) available to treat HIV infections. There is no known cure for HIV or AIDS but the drugs help suppress the virus. By suppressing the amount of virus in the body, people can lead longer and healthier lives. Even though their virus levels may be low they can still spread the virus to others.^[60]



Unscreened

There are many species of bacteria, protozoa, fungi, and viruses, many which remain undocumented or poorly studied with regards to sexual transmission. Despite that the above include what are generally known as STIs, sexually transmission of microbes is far from limited to the above list. Since the sexual route of transmission is not considered common, and/or the microbe itself is not implicated in a major research study on disease, the following pathogens are simply not screened for in sexual health clinics. Some of these microbes are known to be sexually transmittable.

Microbes known to be sexually transmissible (but not generally considered STDs/STIs) include:

- Marburg virus - Virus in semen for 7 weeks after clinical recovery.^[61]
- HTLV (both types 1 and 2) - Sexually transmissible, consumption of breast milk breastfeeding, and once mistaken as a HIV, risk of leukemia.^[62]

Pathophysiology

Many STIs are (more easily) transmitted through the mucous membranes of the penis, vulva, rectum, urinary tract and (less often—depending on type of infection) the mouth, throat, respiratory tract and eyes.^[63] The visible membrane covering the head of the penis is a mucous membrane, though it produces no mucus (similar to the lips of the mouth). Mucous membranes differ from skin in that they allow certain pathogens into the body.^[64] The amount of contact with infective sources which causes infection varies with each pathogen but in all cases a disease may result from even light contact from fluid carriers like venereal fluids onto a mucous membrane.

This is one reason that the probability of transmitting many infections is far higher from sex than by more casual means of transmission, such as non-sexual contact—touching, hugging, shaking hands—but it is not the only reason. Although mucous membranes exist in the mouth as in the genitals, many STIs seem to be easier to transmit through oral sex than through deep kissing. According to a safe sex chart, many infections that are easily transmitted from the mouth to the genitals or from the genitals to the mouth are much harder to transmit from one mouth to another.^[65] With HIV, genital fluids happen to contain much more of the pathogen than saliva. Some infections labeled as STIs can be transmitted by direct skin contact. Herpes simplex and HPV are both examples. KSHV, on the other hand, may be transmitted by deep-kissing but also when saliva is used as a sexual lubricant.

Depending on the STI, a person may still be able to spread the infection if no signs of disease are present. For example, a person is much more likely to spread herpes infection when blisters are present than when they are absent. However, a person can spread HIV infection at any time, even if he/she has not developed symptoms of AIDS.

All sexual behaviors that involve contact with the bodily fluids of another person should be considered to contain some risk of transmission of sexually transmitted diseases. Most attention has focused on controlling HIV, which causes AIDS, but each STI presents a different situation.

As may be noted from the name, sexually transmitted diseases are *transmitted* from one person to another by certain sexual activities rather than being actually *caused by* those sexual activities. Bacteria, fungi, protozoa or viruses are still the causative agents. It is not possible to catch any sexually transmitted disease from a sexual activity with a person who is not carrying a disease; conversely, a person who has an STI got it from contact (sexual or otherwise) with someone who had it, or his/her bodily fluids. Some STIs such as HIV can be transmitted from mother to child either during pregnancy or breastfeeding.

Although the likelihood of transmitting various diseases by various sexual activities varies a great deal, in general, all sexual activities between two (or more) people should be considered as being a two-way route for the transmission of STIs, i.e., "giving" or "receiving" are both risky although receiving carries a higher risk.

Healthcare professionals suggest safer sex, such as the use of condoms, as the most reliable way of decreasing the risk of contracting sexually transmitted diseases during sexual activity, but safer sex should by no means be considered an absolute safeguard. The transfer of and exposure to bodily fluids, such as blood transfusions and other blood products, sharing injection needles, needle-stick injuries

(when medical staff are inadvertently jabbed or pricked with needles during medical procedures), sharing tattoo needles, and childbirth are other avenues of transmission. These different means put certain groups, such as medical workers, and haemophiliacs and drug users, particularly at risk.

Recent epidemiological studies have investigated the networks that are defined by sexual relationships between individuals, and discovered that the properties of sexual networks are crucial to the spread of sexually transmitted diseases. In particular, assortative mixing between people with large numbers of sexual partners seems to be an important factor.

It is possible to be an asymptomatic carrier of sexually transmitted diseases. In particular, sexually transmitted diseases in women often cause the serious condition of pelvic inflammatory disease.

Prevention

Prevention is key in addressing incurable STIs, such as HIV and herpes. Sexual health clinics promote the use of condoms and provide outreach for at-risk communities.

The most effective way to prevent sexual transmission of STIs is to avoid contact of body parts or fluids which can lead to transfer with an infected partner. Not all sexual activities involve contact: cybersex, phonesex or masturbation from a distance are methods of avoiding contact. Proper use of condoms reduces contact and risk. Although a condom is effective in limiting exposure, some disease transmission may occur even with a condom.^[66]

Both partners should get tested for STIs before initiating sexual contact, or before resuming contact if a partner engaged in contact with someone else. Many infections are not detectable immediately after exposure, so enough time must be allowed between possible exposures and testing for the tests to be accurate. Certain STIs, particularly certain persistent viruses like HPV, may be impossible to detect with current medical procedures.

Many diseases that establish permanent infections can so occupy the immune system that other diseases become more easily transmitted. The innate immune system led by defensins against HIV can prevent transmission of HIV when viral counts are very low, but if busy with other viruses or overwhelmed, HIV can establish itself. Certain viral STI's also greatly increase the risk of death for HIV infected patients.

Strategies to increase testing for HIV and STIs have proved to be successful. Some treatment facilities utilize in-home test kits and have the person return the test for follow-up. Other facilities strongly encourage that those previously infected return to ensure that the infection has been eliminated. Novel strategies to foster re-testing have been the use of text messaging and email as reminders. These types of reminders are now used in addition to phone calls and letters.^[67]



San Francisco City Clinic a municipal STI testing center in San Francisco.

Vaccines

Vaccines are available that protect against some viral STIs, such as Hepatitis A, Hepatitis B, and some types of HPV. Vaccination before initiation of sexual contact is advised to assure maximal protection. The development of vaccines to protect against gonorrhea is ongoing.^[68]

Condoms

Condoms and female condoms only provide protection when used properly as a barrier, and only to and from the area that it covers. Uncovered areas are still susceptible to many STIs.

In the case of HIV, sexual transmission routes almost always involve the penis, as HIV cannot spread through unbroken skin; therefore, properly shielding the penis with a properly worn condom from the vagina or anus effectively stops HIV transmission. An infected fluid to broken skin borne direct transmission of HIV would not be considered "sexually transmitted", but can still theoretically occur during sexual contact. This can be avoided simply by not engaging in sexual contact when presenting open, bleeding wounds.

Other STIs, even viral infections, can be prevented with the use of latex, polyurethane or polyisoprene condoms as a barrier. Some microorganisms and viruses are small enough to pass through the pores in natural skin condoms, but are still too large to pass through latex or synthetic condoms.

Proper male condom usage entails:

- Not putting the condom on too tight at the end, and leaving 1.5 cm (3/4 inch) room at the tip for ejaculation. Putting the condom on snug can and often does lead to failure.
- Wearing a condom too loose can defeat the barrier.
- Avoiding inverting or spilling a condom once worn, whether it has ejaculate in it or not.
- If a user attempts to unroll the condom, but realizes they have it on the wrong side, then this condom should be discarded.
- Being careful with the condom if handling it with long nails.
- Avoiding the use of oil-based lubricants (or anything with oil in it) with latex condoms, as oil can eat holes into them.
- Using flavored condoms for oral sex only, as the sugar in the flavoring can lead to yeast infections if used to penetrate.

In order to best protect oneself and the partner from STIs, the old condom and its contents should be assumed to be infectious. Therefore, the old condom must be properly disposed of. A new condom should be used for each act of intercourse, as multiple usage increases the chance of breakage, defeating the effectiveness as a barrier.

Nonoxynol-9

Researchers had hoped that nonoxynol-9, a vaginal microbicide would help decrease STI risk. Trials, however, have found it ineffective^[69] and it may put women at a higher risk of HIV infection.^[70]

Screening

Sexually active women under the age of 25 and those over 25 with risk should be screened for chlamydia and gonorrhoea yearly.^[71] After being treated for gonorrhoea all people should be re tested for the disease after three months.^[71]

Nucleic acid amplification tests are the recommended method of diagnosis for gonorrhoea and chlamydia.^[72] This can be done on either urine in both men and women, vaginal or cervical swabs in women, or urethral swabs in men.^[72]

Diagnosis

Testing may be for a single infection, or consist of a number of tests for a range of STIs, including tests for syphilis, trichomonas, gonorrhoea, chlamydia, herpes, hepatitis and HIV. No procedure tests for all infectious agents.

STI tests may be used for a number of reasons:

- as a diagnostic test to determine the cause of symptoms or illness
- as a screening test to detect asymptomatic or presymptomatic infections
- as a check that prospective sexual partners are free of disease before they engage in sex without safer sex precautions (for example, when starting a long term mutually monogamous sexual relationship, in fluid bonding, or for procreation).
- as a check prior to or during pregnancy, to prevent harm to the baby
- as a check after birth, to check that the baby has not caught an STI from the mother
- to prevent the use of infected donated blood or organs
- as part of the process of contact tracing from a known infected individual
- as part of mass epidemiological surveillance

Early identification and treatment results in less chance to spread disease, and for some conditions may improve the outcomes of treatment. There is often a window period after initial infection during which an STI test will be negative. During this period, the infection may be transmissible. The duration of this period varies depending on the infection and the test. Diagnosis may also be delayed by reluctance of the infected person to seek a medical professional. One report indicated that people turn to the Internet rather than to a medical professional for information on STIs to a higher degree than for other sexual problems.^[73]

Management



World War II US Army poster warning of venereal disease

High-risk exposure such as that which occurs in rape cases may be treated preventatively using antibiotic combinations such as azithromycin, cefixime, and metronidazole.



An option for treating partners of patients (index cases) diagnosed with chlamydia or gonorrhea is patient-delivered partner therapy, which is the clinical practice of treating the sex partners of index cases by providing prescriptions or medications to the patient to take to his/her partner without the health care provider first examining the partner.^[74]

Epidemiology

STD incidence rates remain high in most of the world, despite diagnostic and therapeutic advances that can rapidly render patients with many STDs noninfectious and cure most. In many cultures, changing sexual morals and oral contraceptive use have eliminated traditional sexual restraints, especially for women, and both physicians and patients have difficulty dealing openly and candidly with sexual issues. Additionally, development and spread of drug-resistant bacteria (e.g., penicillin-resistant gonococci) makes some STDs harder to cure. The effect of travel is most dramatically illustrated by the rapid spread of the AIDS virus (HIV-1) from Africa to Europe and the Americas in the late 1970s.^[76]

Commonly reported prevalences of STIs among sexually active adolescent girls both with and without lower genital tract symptoms include chlamydia (10–25%), gonorrhea (3–18%), syphilis (0–3%), *Trichomonas vaginalis* (8–16%), and herpes simplex virus (2–12%). Among adolescent boys with no symptoms of urethritis, isolation rates include chlamydia (9–11%) and gonorrhea (2–3%). A 2008 CDC study found that 25–40% of U.S. teenage girls has a sexually transmitted disease.^{[77][78]}

AIDS is among the leading causes of death in present-day Sub-Saharan Africa.^[79] HIV/AIDS is transmitted primarily via unprotected sexual intercourse. More than 1.1 million persons are living with HIV/AIDS in the United States,^[80] and it disproportionately impacts African Americans.^[81] Hepatitis B is also considered a sexually transmitted disease because it can be spread through sexual contact.^[82] The highest rates



are found in Asia and Africa and lower rates are in the Americas and Europe.^[83] Approximately two billion people worldwide have been infected with the hepatitis B virus.^[84]

History

The first well-recorded European outbreak of what is now known as syphilis occurred in 1494 when it broke out among French troops besieging Naples in the Italian War of 1494–98.^[85] The disease may have originated from the Columbian Exchange.^[86] From Naples, the disease swept across Europe, killing more than five million people.^[87] As Jared Diamond describes it, "[W]hen syphilis was first definitely recorded in Europe in 1495, its pustules often covered the body from the head to the knees, caused flesh to fall from people's faces, and led to death within a few months," rendering it far more fatal than it is today. Diamond concludes, "[B]y 1546, the disease had evolved into the disease with the symptoms so well known to us today."^[88] Gonorrhoeae is recorded at least up to 700 years ago and associated with a district in Paris formerly known as "Le Clapiers". This is where the prostitutes were to be found at that time.^[68]


Prior to the invention of modern medicines, sexually transmitted diseases were generally incurable, and treatment was limited to treating the symptoms of the disease. The first voluntary hospital for venereal diseases was founded in 1746 at London Lock

Hospital.^[89] Treatment was not always voluntary: in the second half of the 19th century, the Contagious Diseases Acts were used to arrest suspected prostitutes. In 1924, a number of states concluded the Brussels Agreement, whereby states agreed to provide free or low-cost medical treatment at ports for merchant seamen with venereal diseases.

The first effective treatment for a sexually transmitted disease was salvarsan, a treatment for syphilis. With the discovery of antibiotics, a large number of sexually transmitted diseases became easily curable, and this, combined with effective public health campaigns against STDs, led to a public perception during the 1960s and 1970s that they have ceased to be a serious medical threat.

During this period, the importance of contact tracing in treating STIs was recognized. By tracing the sexual partners of infected individuals, testing them for infection, treating the infected and tracing their contacts in turn, STI clinics could effectively suppress infections in the general population.

In the 1980s, first genital herpes and then AIDS emerged into the public consciousness as sexually transmitted diseases that could not be cured by modern medicine. AIDS in particular has a long asymptomatic period—during which time HIV (the human immunodeficiency virus, which causes AIDS) can replicate and the disease can be transmitted to others—followed by a symptomatic period, which leads rapidly to death unless treated. HIV/AIDS entered the United States from Haiti in about



1930s Works Progress
Administration poster



U.S. propaganda poster targeted at World War II servicemen appealed to their patriotism in urging them to protect themselves. The text at the bottom of the poster reads, "You can't beat the Axis if you get VD."

1969.^[90] Recognition that AIDS threatened a global pandemic led to public information campaigns and the development of treatments that allow AIDS to be managed by suppressing the replication of HIV for as long as possible. Contact tracing continues to be an important measure, even when diseases are incurable, as it helps to contain infection.

Other names or slang terms

In the 1996 sexual harassment case filed against Mitsubishi Motor Manufacturing of America by the Equal Employment Opportunity Commission (EEOC), a number of male employees referred to sexually transmitted infections as “souvenirs”.^[91]




Plants

- *Microbotryum violaceum* is an STI in a plant caused by a smut fungus and infects the plant species *Silene latifolia*. This smut fungus is spread by pollinators, where pollen is the male gamete in plants, which is why *M. violaceum* is referred to as an STI. It also sterilizes the plants that it infects, regardless of sex.^[92]

References

1. "Sexually transmitted infections (STIs) Fact sheet N°110". *who.int*. November 2013. Retrieved 30 November 2014.
2. Murray PR, Rosenthal KS, Pfaller MA (2013). *Medical microbiology* (7th ed.). St. Louis, Mo.: Mosby. p. 418. ISBN 9780323086929.

3. Goering, Richard V. (2012). *Mims' medical microbiology*. (5th ed.). Edinburgh: Saunders. p. 245. ISBN 9780723436010.
4. "How You Can Prevent Sexually Transmitted Diseases". *cdc.gov*. November 5, 2013. Retrieved 5 December 2014.
5. Centers for Disease Control and Prevention, (CDC) (10 August 2012). "Update to CDC's Sexually transmitted diseases treatment guidelines, 2010: oral cephalosporins no longer a recommended treatment for gonococcal infections.". *MMWR. Morbidity and mortality weekly report*. **61** (31): 590–4. PMID 22874837.
6. GBD 2013 Mortality and Causes of Death, Collaborators (17 December 2014). "Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013.". *Lancet*. **385** (9963): 117–71. doi:10.1016/S0140-6736(14)61682-2. PMC 4340604 . PMID 25530442.
7. "STD Trends in the United States: 2010 National Data for Gonorrhea, Chlamydia, and Syphilis". Centers for Disease Control and Prevention. Retrieved 15 September 2012.
8. Gerd Gross, Stephen K. Tyring (2011). *Sexually transmitted infections and sexually transmitted diseases*. Heidelberg: Springer Verlag. p. 20. ISBN 9783642146633.
9. Organization, World Health (2003). *Guidelines for the management of sexually transmitted infections* (PDF). Geneva: World Health Organization. p. vi. ISBN 9241546263.
10. "Venereal". dictionary.reference.com. Retrieved June 18, 2013.
11. "Sexually transmitted diseases (STDs)?" PLWHA/National AIDS Resource Center. Retrieved March 25, 2013.
12. K. Madhav Naidu. "Epidemiology and Management". *Community Health Nursing*. Gyan Publishing House (2010). p. 248.
13. "Male STI check-up video". Channel 4. 2008. Retrieved 2009-01-22.
14. Hoffman, Barbara (2012). *Williams gynecology*. New York: McGraw-Hill Medical. ISBN 9780071716727.
15. Sarah Edwards, Chris Carn; Carne (1998). "Oral sex and the transmission of non-viral STIs" (PDF). *Sex Transm Infect*. **74** (1): 95–100. doi:10.1136/sti.74.2.95.
16. Maura Gillisons (2007). "HPV Infection Linked to Throat Cancers". *Johns Hopkins Medicine*.
17. Hoare A (2010). models of HIV epidemics in Australia and Southeast Asia (<http://unsworks.unsw.edu.au/fapi/datastream/unsworks:9027/SOURCE02>)
18. Australasian contact tracing manual. Specific infections where contact tracing is generally recommended (<http://ctm.ashm.org.au/Default.asp?PublicationID=6&ParentSectionID=694&SectionID=670>)
19. Varghese B, Maher JE, Peterman TA, Branson BM, Steketee RW (2002). "Reducing the risk of sexual HIV transmission: quantifying the per-act risk for HIV on the basis of choice of partner, sex act, and condom use". *Sex. Transm. Dis*. **29** (1): 38–43. doi:10.1097/00007435-200201000-00007. PMID 11773877.
20. Holmes, King; et al. (1970). "An estimate of the risk of men acquiring gonorrhea by sexual contact with infected females". **91** (2): 170–174.
21. Mahiane, Séverin-Guy; et al. (January 2009). "Transmission probabilities of HIV and herpes simplex virus type 2, effect of male circumcision and interaction: a longitudinal study in a township of South Africa". *AIDS*. **23** (3): 377–383. doi:10.1097/QAD.0b013e32831c5497. PMC 2831044 . PMID 19198042.
22. Burchell A; et al. (March 2006). "Modeling the Sexual Transmissibility of Human Papillomavirus Infection using Stochastic Computer Simulation and Empirical Data from a Cohort Study of Young Women in Montreal, Canada". *American Journal of Epidemiology*. **169** (3): 534–543. doi:10.1093/aje/kwj077. PMID 16421235.
23. Platt, Richard; et al. (1983). "Risk of Acquiring Gonorrhea and Prevalence of Abnormal Adnexal Findings Among Women Recently Exposed to Gonorrhea". *JAMA*. **250** (23): 3205–3209. doi:10.1001/jama.250.23.3205. PMID 6417362.
24. Department of Public Health, City & County of San Francisco (2011).STD Risks Chart (<http://www.sfcityclinic.org/stdbasics/stdchart.asp>)
25. Jin F; et al. (March 2010). "Per-contact probability of HIV transmission in homosexual men in Sydney in the era of HAART". *AIDS*. **24** (6): 907–913. doi:10.1097/QAD.0b013e3283372d90. PMC 2852627 . PMID 20139750.
26. Bryan C (2011)INFECTIOUS DISEASE CHAPTER EIGHT SEXUALLY TRANSMITTED DISEASES (<http://pathmicro.med.sc.edu/infectious%20disease/sexually%20transmitted%20diseases.htm>)
27. Richard Pearson (2007). "Pinworm Infection". *Merck Manual Home Health Handbook*.

28. Caini, Saverio; Gandini, Sara; Dudas, Maria; Bremer, Viviane; Severi, Ettore; Gherasim, Alin (2014). "Sexually transmitted infections and prostate cancer risk: A systematic review and meta-analysis". *Cancer Epidemiology*. **38** (4): 329–338. doi:10.1016/j.canep.2014.06.002. ISSN 1877-7821. PMID 24986642.
29. Ljubin-Sternak, Sunčanica; Meštrović, Tomislav (2014). "Chlamydia trachomatis and Genital Mycoplasmas: Pathogens with an Impact on Human Reproductive Health". *Journal of Pathogens*. **2014**: 1–15. doi:10.1155/2014/183167. ISSN 2090-3057. PMC 4295611 . PMID 25614838.
30. Schlicht, M. J.; Lovrich, S. D.; Sartin, J. S.; Karpinsky, P.; Callister, S. M.; Agger, W. A. (2004). "High Prevalence of Genital Mycoplasmas among Sexually Active Young Adults with Urethritis or Cervicitis Symptoms in La Crosse, Wisconsin". *Journal of Clinical Microbiology*. **42** (10): 4636–4640. doi:10.1128/JCM.42.10.4636-4640.2004. ISSN 0095-1137.
31. McIver, C. J.; Rismanto, N.; Smith, C.; Naing, Z. W.; Rayner, B.; Lusk, M. J.; Konecny, P.; White, P. A.; Rawlinson, W. D. (2009). "Multiplex PCR Testing Detection of Higher-than-Expected Rates of Cervical Mycoplasma, Ureaplasma, and Trichomonas and Viral Agent Infections in Sexually Active Australian Women". *Journal of Clinical Microbiology*. **47** (5): 1358–1363. doi:10.1128/JCM.01873-08. ISSN 0095-1137.
32. Clark, Natalie; Tal, Reshef; Sharma, Harsha; Segars, James (2014). "Microbiota and Pelvic Inflammatory Disease". *Seminars in Reproductive Medicine*. **32** (01): 043–049. doi:10.1055/s-0033-1361822. ISSN 1526-8004. PMC 4148456 . PMID 24390920.
33. Larsen, Bryan; Hwang, Joseph (2010). "Mycoplasma, Ureaplasma, and Adverse Pregnancy Outcomes: A Fresh Look". *Infectious Diseases in Obstetrics and Gynecology*. **2010**: 1–7. doi:10.1155/2010/521921. ISSN 1064-7449.
34. "Giardia, Epidemiology & Risk Factors". Center For Disease Control. July 13, 2012. Retrieved 2015-07-03.
35. "Hepatitis A, Division of Viral Hepatitis". Center For Disease Control. May 31, 2015. Retrieved 2015-07-03.
36. "Shigella Infections among Gay & Bisexual Men". Center For Disease Control. April 23, 2015. Retrieved 2015-07-03.
37. Ljubin-Sternak, Sunčanica; Mestrovic, Tomislav (2014). "Review: Chlamydia trachomatis and Genital Mycoplasmas: Pathogens with an Impact on Human Reproductive Health". *Journal of Pathogens*. **2014** (183167): 1–15. doi:10.1155/2014/183167. PMC 4295611 . PMID 25614838.
38. Zarei, Omid; Rezaia, Simin; Mousavi, Atefeh (2013). "Mycoplasma genitalium and Cancer: A Brief Review". *Asian Pacific Journal of Cancer Prevention*. **14** (6): 3425–3428. doi:10.7314/APJCP.2013.14.6.3425. ISSN 1513-7368.
39. Manchester, Marianne; McGowin, Chris L.; Anderson-Smits, Colin (2011). "Mycoplasma genitalium: An Emerging Cause of Sexually Transmitted Disease in Women". *PLoS Pathogens*. **7** (5): e1001324. doi:10.1371/journal.ppat.1001324. ISSN 1553-7374.
40. Taylor-Robinson D, Clin Infect Dis. 1996 Oct;23(4):671-82; quiz 683-4. Infections due to species of Mycoplasma and Ureaplasma: an update.
41. Workowski K, Berman S (2006). "Sexually transmitted diseases treatment guidelines, 2006." (PDF). *MMWR Recomm Rep*. **55** (RR-11): 1–94. PMID 16888612.
42. Wu J, Chen C, Sheen I, Lee S, Tzeng H, Choo K (1995). "Evidence of transmission of hepatitis D virus to spouses from sequence analysis of the viral genome". *Hepatology*. **22** (6): 1656–60. doi:10.1002/hep.1840220607. PMID 7489970.
43. Farci P (2003). "Delta hepatitis: an update". *J Hepatol*. **39** (Suppl 1): S212–9. doi:10.1016/S0168-8278(03)00331-3. PMID 14708706.
44. Shukla N, Poles M (2004). "Hepatitis B virus infection: co-infection with hepatitis C virus, hepatitis D virus, and human immunodeficiency virus". *Clin Liver Dis*. **8** (2): 445–60, viii. doi:10.1016/j.cld.2004.02.005. PMID 15481349.
45. King, B. (2009). *Human Sexuality Today* (Sixth ed.). Upper Saddle River: Pearson Education, Inc.
46. "Chlamydia Infections: MedlinePlus". Nlm.nih.gov. Retrieved 2013-06-30.
47. <http://www.webmd.com/genital-herpes/guide/genital-herpes-basics>
48. "Herpes". Avert.org. Retrieved 2013-06-30.
49. "Human Papillomavirus (HPV) | Overview". FamilyDoctor.org. 2010-12-01. Retrieved 2013-06-30.
50. "STD Facts - Human papillomavirus (HPV)". Cdc.gov. Retrieved 2013-06-30.
51. "Gonorrhea - National Library of Medicine - PubMed Health". Ncbi.nlm.nih.gov. Retrieved 2013-06-30.
52. "STD Facts –Syphilis". Cdc.gov. Retrieved 2013-02-18.
53. "Syphilis". Who.int. Retrieved 2013-02-18.

54. Clement, Meredith E.; Okeke, N. Lance; Hicks, Charles B. (2014). "Treatment of Syphilis". *JAMA*. **312** (18): 1905. doi:10.1001/jama.2014.13259. ISSN 0098-7484.
55. "STD Facts – Trichomoniasis". Cdc.gov. Retrieved 2013-02-18.
56. "Trichomoniasis: MedlinePlus". Nlm.nih.gov. Retrieved 2013-02-18.
57. "Trichomoniasis – NHS Choices". Nhs.uk. 27 February 2012. Retrieved 2013-02-18.
58. "Hiv/Aids". Mayo Clinic.com. 2012-08-11. Retrieved 2013-06-30.
59. "AIDS". Avert.org. Retrieved 2013-06-30.
60. "HIV/AIDS Treatment". Niaid.nih.gov. 2009-06-03. Retrieved 2013-06-30.
61. "Marburg virus disease: origins, reservoirs, transmission and guidelines,". (Great Britain] GOV.UK. 5 September 2014. Retrieved 2015-07-03.
62. Verdonck, K.; González, E.; Van Dooren, S.; Vandamme, A. M.; Vanham, G.; Gotuzzo, E. (April 2007). "Human T-lymphotropic virus 1: Recent knowledge about an ancient infection". *The Lancet Infectious Diseases*. **7** (4): 266–281. doi:10.1016/S1473-3099(07)70081-6. PMID 17376384.
63. K. Madhav Naidu. *Community Health Nursing*, Gen Next Publications, 2009, p.248
64. Virus Pathogenesis (<http://www.microbiologybytes.com/virology/VirPath.html>), Microbiology Bytes
65. Safe Sex Chart (<http://www.tinybubbles.com/safesexchart.html>), violet blue: open source sex
66. Villhauer, Tanya (2005-05-20). "Condoms Preventing HPV?". University of Iowa Student Health Service/Health Iowa. Retrieved 2009-07-26.
67. Desai, Monica; Woodhall, Sarah C; Nardone, Anthony; Burns, Fiona; Mercey, Danielle; Gilson, Richard (2015). "Active recall to increase HIV and STI testing: a systematic review". *Sexually Transmitted Infections: sextrans-2014-051930*. doi:10.1136/sextrans-2014-051930. ISSN 1368-4973.
68. Baarda, Benjamin I.; Sikora, Aleksandra E. (2015). "Proteomics of Neisseria gonorrhoeae: the treasure hunt for countermeasures against an old disease". *Frontiers in Microbiology*. **6**. doi:10.3389/fmicb.2015.01190. ISSN 1664-302X; Access provided by the University of Pittsburgh.
69. Wilkinson D, Ramjee G, Tholandi M, Rutherford G (2002). Wilkinson D, ed. "Nonoxynol-9 for preventing vaginal acquisition of sexually transmitted infections by women from men". *Cochrane Database Syst Rev* (4): CD003939. doi:10.1002/14651858.CD003939. PMID 12519623.
70. Williams, M. *Healthy Choices for Fertility Control*. CreateSpace, Scotts Valley, CA, 2009. ISBN 1-4486-6472-1
71. Gavin L, Moskosky S, Carter M, Curtis K, Glass E, Godfrey E, Marcell A, Mautone-Smith N, Pazol K, Tepper N, Zapata L (Apr 25, 2014). Division of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion, CDC. "Providing Quality Family Planning Services: Recommendations of CDC and the U.S. Office of Population Affairs". *MMWR. Recommendations and reports : Morbidity and mortality weekly report. Recommendations and reports / Centers for Disease Control*. **63** (RR-04): 1–54. PMID 24759690.
72. "Screening for Chlamydia and Gonorrhea: U.S. Preventive Services Task Force Recommendation Statement". *USPSTF*. Retrieved 29 April 2014.
73. Quilliam Susan (2011). "The Cringe Report" ". *J Fam Plann Reprod Health Care*. **37** (2): 110–112.
74. Expedited Partner Therapy in the Management of Sexually Transmitted Diseases (2 February 2006) (<http://www.cdc.gov/std/treatment/EPTFinalReport2006.pdf>) U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES PUBLIC HEALTH SERVICE. Centers for Disease Control and Prevention National Center for HIV, STD, and TB Prevention
75. "WHO Disease and injury country estimates". *World Health Organization*. 2004. Retrieved Nov 11, 2009.
76. Mary-Ann Shafer; Anna-Barbara Moscicki (2006). "Sexually Transmitted Infections, 2006": 1–8.
77. "Sex Infections Found in Quarter of Teenage Girls (http://www.nytimes.com/2008/03/12/science/12std.html?_r=1)". The New York Times. March 12, 2008.
78. "CDC study says at least 1 in 4 teen girls has a sexually transmitted disease; HPV most common (<http://newsok.com/article/3214748/1205233095>)". The Oklahoman. March 11, 2008.
79. UNAIDS, WHO (December 2007). "2007 AIDS epidemic update" (PDF). Retrieved 2008-03-12.
80. "HIV in the United States: At A Glance (<http://www.cdc.gov/hiv/statistics/basics/ataglance.html>)". Centers for Disease Control and Prevention.
81. "AIDS In Black America: A Public Health Crisis (<http://www.npr.org/2012/07/05/156292172/aids-in-black-america-a-public-health-crisis>)". NPR. July 5, 2012.
82. "Hepatitis: Type B (caused by hepatitis B virus) (<http://www.health.state.mn.us/divs/idepc/diseases/hepb/hepbstd.html>)". Minnesota Department of Health.

83. "Hepatitis B (http://www.fda.gov/ohrms/dockets/ac/02/slides/3885S2_02_Hoofnagle/tsld002.htm)". U.S. Food and Drug Administration.
84. "World Hepatitis Day 2012". WHO. 2012.
85. Oriel, J.D. (1994). *The Scars of Venus: A History of Venereology*. London: Springer-Verlag. ISBN 3-540-19844-X.
86. "Columbus May Have Brought Syphilis to Europe (<http://www.livescience.com/history/080114-syphilis-columbus.html>)". LiveScience. January 15, 2008.
87. CBC News Staff (January 2008). "Study traces origins of syphilis in Europe to New World". Retrieved 2014-02-21.
88. Diamond, Jared (1997). *Guns, Germs and Steel*. New York: W.W. Norton. p. 210. ISBN 84-8306-667-X.
89. Archives in London and the M25 area (AIM25 (<http://www.aim25.ac.uk/index.stm>)) London Lock Hospital records (http://www.aim25.ac.uk/cgi-bin/search2?coll_id=859&inst_id=9)
90. Gilbert MT, Rambaut A, Wlasiuk G, Spira TJ, Pitchenik AE, Worobey M (November 2007). "The emergence of HIV/AIDS in the Americas and beyond". *Proc. Natl. Acad. Sci. U.S.A.* **104** (47): 18566–70. doi:10.1073/pnas.0705329104. PMC 2141817. PMID 17978186. Retrieved March 20, 2010.
91. "Equal Employment Opportunity Commission v. Mitsubishi Motor Manufacturing of America Inc." (<http://www.contilaw.com/pleadings/EEOCvMitsubishsi.htm>), Adam J. Conti, LLC, Filing 96-1192, September 15, 1997
92. Kaltz, Oliver; Shykoff, Jacqui A. (1 Feb 2001). "Male and Female *Silene latifolia* Plants Differ in Per-Contact Risk of Infection by a Sexually Transmitted Disease". *Journal of Ecology*. British Ecological Society. **89** (1): 99–109.

Further reading

- Workowski, KA; Berman, S; Centers for Disease Control and Prevention, (CDC) (17 December 2010). "Sexually transmitted diseases treatment guidelines, 2010.". *MMWR. Recommendations and reports : Morbidity and mortality weekly report. Recommendations and reports / Centers for Disease Control*. **59** (RR-12): 1–110. PMID 21160459.
- Aral, Sevgi O (2008). *Behavioral Interventions for Prevention and Control of Sexually Transmitted Diseases*. : Springer Singapore Pte. Limited. ISBN 978-0-387-85768-8
- Faro, Sebastian (2003). *Sexually transmitted diseases in women*. Lippincott Williams & Wilkins. ISBN 0-397-51303-8
- Ford, Carol A; Elizabeth Shimer Bowers (2009). *Living with Sexually Transmitted Diseases*. Facts On File. ISBN 978-0-8160-7672-7
- Sehgal, Virendra N (2003). *Sexually Transmitted Diseases* (4th ed.). Jaypee Bros. Medical Publishers. ISBN 81-8061-105-1
- Shoquist, Jennifer; Diane Stafford (2003). *The encyclopedia of sexually transmitted diseases*. Facts On File. ISBN 0-8160-4881-9

External links

- Sexually transmitted infection



(https://www.dmoz.org/Health/Reproductive_Health/Sexually_Transmitted_Diseases/) at DMOZ

- CDC Sexually Transmitted Diseases Treatment Guidelines, 2010 (<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5912a1.htm>)
- STD photo library (<http://www.dermnet.com/moduleIndex.cfm?moduleID=16>) at Dermnet

- UNFPA: Breaking the Cycle of Sexually Transmitted Infections (<http://www.unfpa.org/rh/stis.htm>) at UNFPA
- STDs In Color: Sexually Transmitted Disease Facts and Photos (<http://www.stdsincolor.com/>)
- [1] (<http://wonder.cdc.gov/std.html>): Sexually transmitted diseases in the U.S.

Retrieved from "https://en.wikipedia.org/w/index.php?title=Sexually_transmitted_infection&oldid=749794644"

Categories: [Human sexuality](#) | [Sexually transmitted diseases and infections](#) | [Sexual health](#) | [Midwifery](#)

- This page was last modified on 16 November 2016, at 05:38.
- Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.